[45] **Jul. 18, 1978**

[54]	PROCESS FOR THE CROSS-LINKING OF PROTEINS		
[75]	Inventors:	Menachem Rubinstein, Kiryat Ono; Shulamit Simon, Ramat-Gan, both of Israel; Rene Bloch, deceased late of Savion, Israel, by Rachel Bloch, administratrix	
[73]	Assignee:	Research Products Rehovot Ltd., Rehovot, Israel	
[21]	Appl. No.:	695,030	
[22]	Filed:	Jun. 11, 1976	
[30]	Foreign Application Priority Data		
Jun. 12, 1975 [IL] Israel			
[51] [52]	1] Int. Cl. ² C07G 7/02; C07G 7/00 2] U.S. Cl 195/63; 195/68; 195/DIG. 11; 260/112 R; 260/122		
[58]	[58] Field of Search		
[56]		References Cited	
U.S. PATENT DOCUMENTS			
	96,634 3/19 99,267 2/19		
OTHER PUBLICATIONS			
Wold in Hirs (Editor) Methods in Enzymology, vol. XI (1967), pp. 617-640.			

Primary Examiner—Lionel M. Shapiro

Attorney, Agent, or Firm-Millen & White

[57] ABSTRACT

The invention relates to a process for the cross-linking of proteins wherein the proteins to be cross-linked are reacted with a cross-linking agent selected from the group consisting of:

(a) water soluble poly(ethylene oxide) derivatives of the general formula I

$$(Z)_m - Y - (CH_2CH_2O)_n CH_2CH_2 - Y - (Z)_m$$
 (I)

wherein both n and m are at least 1; Y is a covalent bond or is an —R— or —RO— radical in which the oxygen is bound to the poly(ethylene oxide) and R is selected from the group consisting of methylene, ethylene, propylene, o—, m— and p-phenylene, o—, m— and p-phenylene carbamate optionally substituted by one or more alkyl, aryl, halo, nitro, oxo, carboxy, hydroxy, thio, sulfonate and phosphate groups; and Z is a reactive group selected from the groups consisting of haloisocyanato-, isothiocyanato-, tosylate, acyl halides, acyl azides, aryl diazonium salts, acyl imidoester salts, activated esters of acyl residues and 2,4-dichlorotriazines;

(b) activated esters of di- and poly-carboxylic acids wherein the acids and the alcohol moieties thereof are water soluble.

31 Claims, No Drawings